

Corrections and clarifications in *BNF for Children*

Date posted	Edition	Versions affected	Clarification or correction		Comment
			Location	Detail	
17 October 2007	2007	Book & Digital	Page 74, column 1 Section 1.5	<p>The dose of intravenous ciclosporin for the treatment of refractory ulcerative colitis in children is incorrect. The dose by intravenous infusion should read:</p> <p>Child 3–18 years initially 0.5–1 mg/kg twice daily; dose adjusted according to blood-ciclosporin concentration and response</p>	<p>The error will be corrected for the print edition of BNFC 2008</p> <p>Online versions of BNFC 2007 have been corrected. An update to correct this error will be available for other digital versions of BNFC 2007</p>
8 January 2007	2006	Book	Page 750 Top of column 1 and column 2	Column header should read 'Antimalarials' not 'Antihistamines'	The error will be corrected for BNFC 2007

27 November 2006	2006	Book & digital	Page 854, column 1	BNFC2006 shows an incorrect email address for IVAX Pharmaceuticals.	Queries to IVAX should instead be made by post, telephone, or fax
13 November 2006	2006	Book & digital	page 268 right-hand column section 4.8.1	The strength of Nitrazepam oral suspension is incorrect. The preparation should read as follows:	The error will be corrected for BNFC 2007
				Oral suspension, nitrazepam 2.5 mg/5 mL, net price 150 mL = £5.30. Label: 1, 8	
3 November 2006	2006	Book & digital	page 261 right-hand column section 4.8.1	In the paragraph on Administration (under Phenobarbital), the rate of injection is incorrect and the statement on intravenous injection should read:	The error will be corrected for BNFC 2007. Note that the rate of injection shown on p.272 (section 4.8.2) is correct.
				For <i>intravenous injection</i> , dilute to a concentration of 20 mg/mL with Water for Injections; give over 20 minutes (no faster than 1 mg/kg/minute).	

31 October 2006	2006	Book & digital	page 680 section 14.4	<p>The strength of one of the Tuberculin Purified Protein Derivative preparations is incorrect. The preparation entry should read:</p> <p>Injection, heat-treated products of growth and lysis of appropriate <i>Mycobacterium</i> spp. 20 units/mL (2 units/0.1-mL dose) (for routine use), 1.5-mL vial; 100 units/mL (10 units/0.1-mL dose), 1.5-mL vial</p>	The error will be corrected for BNFC 2007
22 May 2006	2005	Book & digital	page 164, column 1 & section 3.1.1	<p>The dose shown under the entry for <i>Oxis[®] Turbohaler[®]</i> should read as follows:</p> <p>Chronic asthma</p> <ul style="list-style-type: none"> By inhalation of powder <p>Child 6–18 years 9 micrograms 1–2 times daily; occasionally up to 36 micrograms daily may be needed (max. single dose 9 micrograms); reassess treatment if additional doses required on more than 2 days a week</p> <p>Relief of bronchospasm</p> <ul style="list-style-type: none"> By inhalation of powder <p>Child 6–18 years 4.5–9 micrograms</p> <p>Prevention of exercise-induced bronchospasm</p> <ul style="list-style-type: none"> By inhalation of powder <p>Child 6–18 years 4.5–9 micrograms before exercise</p>	BNFC 2006 will show the correct doses

1 October 2005	2005	Book (first print only— see Comment)	page 290, column 1	<p>There is an error in the dose of cefalexin for a child aged 12–18 years. The dose should read as follows:</p> <p>Child 12–18 years 500 mg 2–3 times daily, increased to 1–1.5 g 3–4 times daily for severe infection</p>	Digital versions of <i>BNF for Children</i> have been amended to show the correct dose. So too have copies of the book reprinted in October 2005
8 July 2009	2008 2009	Book & digital Book	Page 258, column 1 Page 259, column 1 Section 4.7.2	<p>The route of administration for tramadol for the treatment of postoperative pain in children is incorrect. The dose should be administered by intravenous injection and not by mouth. The dose should read as follows:</p> <ul style="list-style-type: none"> By intravenous injection (over 2–3 minutes) <p>Child 12–18 years 100 mg initially then 50 mg every 10–20 minutes if necessary up to total max. 250 mg (including initial dose) in first hour, <i>then</i> 50-100 mg every 4-6 hours, max. 600 mg daily</p>	The error will be corrected promptly on bnfc.org and for BNFC 2010 print edition